




k	2 - Leading Cause Cause of Death	~	Rate	Deaths
•	es of the heart (heart disease)	\neg	203.3	680,981
	nant Neoplasms (cancer)	=	183.1	613,352
	ents (unintentional injuries)	٦ŀ	66.5	222,698
	ovascular diseases (stroke)	Ħ١	48.6	162,639
Chroni	ic lower respiratory diseases	T)	43.4	145,357
Alzhei	mer's disease		34.0	114,034
Diabet	es mellitus (diabetes)		28.4	95,190
Nephr	itis, nephrosis (kidney disease)		16.5	55,253
Chron	ic liver disease and cirrhosis		15.6	52,222
COVI	D-19		14.9	49,932
Suici	ide (intentional self-harm)		14.7	49,316










8

C	aus	es of	f Death by Age	in U	SA C
Ranks h	igher	for yo	oung,		SA N
lower f	or old	der adı	ults <u>Older Adults 65+ ye</u>	ars ol	AS
			Rank & Cause of Death	Rate	Deaths
<u>Youth 15</u>	<u>~24 ye</u>	<u>ars oias</u>	Diseases of the heart	935.7	551,433
Cause	Pata	Deaths	2. Malignant neoplasms	778.7	461,345
			Cerebrovascular diseases	237.7	140,813
 Accidents 	32.2	14,126	 Chronic lower respiratory diseases 	212.0	125,603
2. Suicide	13.5	5.936	Alzheimer's disease	190.0	112,548
			Accidents (Unintentional injuries)	124.6	73,804
Homicide	13.1	5,745	7. Diabetes mellitus	115.7	68,550
			8. Nephritis & nephrosis	76.3	45,200
33,711	total de	eaths	9. COVID-19	74.4	44,097
	Rate=76.8		10. Parkinson's disease	66.2	39,238
			 Influenza & pneumonia 	62.2	36,859
	-00	100	12. Hypertension	59.3	35,131
1	コンはい	(2 (S) C	13. Septicemia	53.9	31,914
-/	XX M	UNIVOLET	14. Chronic liver disease & cirrhosis	36.9	21,887
_Ω(N W	48341 JA E	15. Nutritional deficiencies	35.8	21,203
Ω6 78	4 Kod 3K7	11 1693 71,	16. Pneumonitis due to solids and liquid	28.5	16,867
A.S.A.	AKSYOKY		17. Benign neoplasms	23.1	13,689
- F 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	*****	RATTINK IT	18. Suicide (Intentional self-harm)	17.6	10,437
	17.6	% of young deaths	s I 0.5% of elderly deaths		2,310,872
	2004 0	ankina .	cauca for vouna (15 28 s	me3	
	cnu ri	ankiny (cause for young (15-24 y	rs)	total deaths




10







12








14


















20







22













26











32

34





Suicidal Behavior Estimates 2023 SAMHSA (2024) study suggests: In the past year: • 856,000 American 1.5 million American adolescents attempted adults attempted suicide (3.3% of suicide (0.6% of adults) adolescents) 3.7 million adults • 1.5 million adolescents made a plan (1.4% of made a plan (5.6% of adults) adolescents) • 12.8 million adults •3.2 million adolescents had serious thoughts had serious thoughts (5.0% of adults) (12.3% of adolescents) **Adults** Adolescents National Survey on Dru





33


Attempted Suicides (Nonfatal Based on 2023's 2.356 million attempted suicides, there is 1 attempt every 13 seconds






























44







46













50








52





























62







64













68







70












74








76

















Sex & Suicio			Suicide Prevention	100			
Youths age	d 15.2	Д 1)	VC	·Cip			
Firearms the lead	ding metho	d for 1	oung Males	S lational Co			
Firearms the leading method for Young Males Young Females used hanging most, then firearms Males Firearms 61.2% 2,915 30.5% 359							
	<u>Male</u>	<u>S</u>	<u>Femal</u>	les 🗓			
Firearms	61.2%	2,915	30.5%	359 g			
$Suffocation \ _{incl.\ hanging,\ strangulation}$							
$Poisoning \ \ (solid \ and \ liquid \ and \ gas)$	5.8%	278	21.9%	258			
All Other Methods	9.3%	445	11.2%	132			
Total Number	r 4,760		1,176				
2023 Data			rading cause of dea young males ages y not equal 100% due to				






Causes of Death by Age in USA Percent of deaths (rate)					
Rank	5-44				
1	Unintentional Injury 82,046	38.6%	(46.9)		
2	Suicide 23,409	11.0%	(13.4)		
3	Malignant Neoplasms 17,107	8.1%	(9.8)	Suicide ranks 2 nd	
4	Homicide 16,585	7.8%	(9.5)	for ages 5 to 44 years	
5	Heart Disease 16.010	7.5%	(9.2)	2 22	

















92








94

































Sex & Suicide Methods: Middle Age - 45-64 years Firearms the leading method for Middle Age Males Middle Age Females used poisoning most often, firearms close second							
	Males		<u>Femal</u>	les 🖁			
Firearms	58.6%	6,964	34.7%	1,244			
$Suffocation \ \ {\it incl.hanging,strangulation}$	24.0%	2,856	20.8%	744			
$Poisoning \ \ (solid \ and \ liquid \ and \ gas)$	9.0%	1,072	35.3%	1,263			
All Other Methods	8.4%	996	9.2%	330			
Total Number	,	e: Totals may	3,581	rounding 10			








Sex & Suicide Methods: Ages - 25-44 years Firearms the leading method for Males Age 25-44 but use of suffocation higher than in adults 45+							
Females Age 25-44 used firearms							
	<u>Male</u>	<u>S</u>	<u>Fema</u>	<u>les</u> 🛚			
Firearms	51.8%	7,009	36.1%	1,243			
Suffocation incl. hanging, strangulation	32.7%	4,422	32.1%	1,104			
$Poisoning \ \ (\text{solid and liquid and gas})$	7.2%	979	22.5%	775			
All Other Methods	8.4%	1,133	9.3%	321			
Total Number	r 13,543		3,443				
	The low use of only in the 25	-44 and		oups			


110































































Suicide Rates by Hispanic Origin, 1990-2023 NH AIAN NH Asia NH Asian = -Hispanic American Indian/Alaska Native highest in recent years














135














141

